

POKE PLAN

For Children Ages 0 - 3 years old

CIRCLE YOUR CHOICES BELOW

I think this
pain reducing tool
with help my child best



Buzzy Bee Ice Pack



Shot Blocker




Sugar Water
(Make at home & bring to office)



Numbing Cream
(Buy OTC & Apply at Home before visit)

I think this
distraction
will help my child best



Look at a Headband



Watch a Pinwheel




Listen to Music



Watch a Video on Your Phone


I think this
position
will make my child feel
safe & comfortable



Your child sits on your lap, with both legs to one side. You wrap both arms around theirs in a comforting hug. This reminds them to keep their arms still, while in a comforting embrace. The child can look at the poke, or choose to look away.



Your child sits on your lap, facing away from you. You wrap both arms around theirs in a comforting hug. You can also wrap your legs around theirs for a full embrace. For bigger kids, you can have them sit on a chair or bed, and straddle them from behind.



Your child sits on your lap, facing you. Their legs straddle and wrap around your waist. You wrap both arms around theirs, for a full embrace, using your underarms and forearms to keep their arms safely contained. This works even for older children.



Your child sits on a table, with you hugging them from behind. You wrap your arms around theirs for a big hug. Use your hugging arms to keep their hands safely contained. For smaller children, you can also use your arms to remind their legs to stay still.

MAKE SURE TO GRAB A STICKER ON YOUR WAY OUT!

RECIPE FOR



- You can use sugar water to reduce your baby's pain. Sugar water is safe for babies, even newborns.
- Make sugar water at home by mixing 1 teaspoon of white sugar with 1 ounce of water.
- Give your baby some sugar water 1 or 2 minutes before the needle.
- Use a dropper (or syringe) to place the sugar water into the side of your baby's mouth between the cheeks and gums. Give your baby one drop at a time and let your baby suck on the sweet taste.
- You may also dip a soother (pacifier) into the sugar water and give it to your baby before, during, and after the needle.

Numbing cream


When, where and how should I apply?

When should I apply?

Approximately 30 min prior to visit. Leave it on until it is time for the poke>


Where should I apply?

Child under 5 years old – Apply to both legs



1. Lay your child down facing you.
2. Draw an imaginary line down the middle of the front of one of your child's legs, from the hip to the knee (line A).
3. Next divide the upper leg into 3 equal sections (lines B & C). Apply the cream or patch to the middle outer section of the leg.

Child older than 5 years – Apply to both arms



How should I apply?

Step 1: *We recommend over the counter LMX numbing cream*

Step 2: Gather supplies – Numbing cream, tegederm or plastic wrap

Step 3: Do not clean skin first (Natural oils help with absorption). Apply a small amount and rub in completely

Step 4: On top of that apply a large glob and DO NOT rub in.

Step 5: Cover with tegederm or plastic wrap

Numbing cream video: https://www.youtube.com/watch?v=YrnSUJWU_30